

ARTYKUŁ POGŁĄDOWY / REVIEW PAPER

Otrzymano/Submitted: 10.06.2019 • Zaakceptowano/Accepted: 20.06.2019

© Akademia Medycyny

What anaesthesiologist should know about the leucocytes and their effect on microcirculation

Krzysztof Kusza¹, Jacek B. Cywinski², Maria Siemionow³

¹ Chair and Department of Anaesthesiology and Intensive Therapy, Poznan University of Medical Sciences, Poznan, Poland

² Department of General Anaesthesiology, Cleveland Clinic, Cleveland, OH, USA

³ Department of Orthopaedics, University of Illinois, Chicago, IL, USA



Abstract

Alterations in microvascular circulation are observed in different pathological conditions, but acute changes in morphotic elements of blood are not well recognised. In conditions like septic shock, severe cardiac failure and others extreme pathophysiologic states, these alteration may influence patient's outcome. Therefore the knowledge of leukocytes behaviour at microcirculatory level in pathophysiologic conditions could be helpful in understanding how different choice of anaesthetic affects microcirculation. *Anestezjologia i Ratownictwo 2019; 13: 122-126.*

Keywords: microcirculation, volatile anaesthetic agents, leucocytes, endothelium, reactive oxygen species, cytokines

the capillaries, depending on how long the deformation process lasts [13-15]. It is unclear what is more important in terms of effect on the microcirculation of the blood flow: deformation time or occlusion time. Mathematical formulas developed by Needham and Hochmuth in 1989 and Fenton in 1985 provided no definite answer to this question. They suggest that the higher the pressure force of the inflowing blood up stream, the shorter the deformation time [16,17]. Adhesion of white cells to the vascular wall is more frequent when the blood flow in the vessel is slowed down [18]. This promotes margination of white cells, which thus become more susceptible to chemotactic substances [19].

It is known that the higher volume and viscosity of white blood cells compared to red blood cells (one white cell is equivalent to 700 red cells) leads to much faster mechanical arrest of blood flow in vessels 5 micrometres in diameter [20,21]. Other *in vivo* studies proved that the increase of resistance in the capillary network and the incidence of lumen occlusion by leukocytes are dependent on the capillary network structure. Variation of pressure values in the capillary network, depending on the number of occlusions, also depends on intervascular canals within the network. A higher number of canals within the capillary network leads to lower variation of pressure values, in consequence increasing the time of vessel occlusion by leukocytes. Reversely, in borderline conditions where no canals between the capillaries are present within the network, the pressure increase will correspond to the pressure value at the capillary network entrance, and the leukocyte occlusion will be very quickly reversed [11,12,21].

Another important feature of the capillary network geometry is the number of branching within the network, as they affect the leukocyte behaviour [22,23]. Experimental models of microcirculation in skeletal muscle showed that significant number of vessels must become occluded before delayed vascular resistance increase is observed in the network [9,15,24]. For instance, a network model comprised of 60 parallel vessels shows no signs of resistance increase until at least five main paths of this network are totally occluded. Above this number, a slight increase in the vascular resistance, not exceeding 9% of the baseline value, was recorded. In the tree-branched vascular network, occlusion of a small number of capillaries causes a high increase in flow resistance since in this structure a much larger number of capillaries remains devoid of blood flow [13,15,25]. There are differences in

capillary network structure, which may significantly affect flow disturbances and change the so-called capillary reserve, depending on the organ being affected or subjected to surgery [12,26].

The effect of white blood cells on the increase in flow resistance is also dependent on changes in blood viscosity. Medications and substances, as well as pathological lesions formed in the system and affecting white cell deformation and viscosity affect the capillary network resistance [13,27].

In addition, occlusion of capillaries by white cells affect further white cell distribution in the capillary network and is directly responsible for the phenomenon of highest flow pathway selection [28]. When one of the vessels is occluded, white blood cells flow is transferred to another capillary network. The more vessels are occluded, the more white cells are dispersed within the entire network [4,5,29].

Postcapillary venules are primary receptacles for blood passing through the capillary network; concurrently it is the last anatomical and functional structure of microcirculation [30]. They play a key role in venous insufficiency of a free skin flap, free muscle flap, composed of multiple tissues or transplanted organs [31,32]. Blockage of flow through some of postcapillary venules results in congestion of an isolated or transferred muscle flap or other tissue, with slowed blood flow through its arteries and is often combined with irreversible changes due to severe abnormalities of metabolism in cells located in the affected area [33,34].

Reduced accumulation of white blood cells in the postcapillary venule directly results in improvement of capillary perfusion, significantly delays development of endothelial edema that eventually results in complete blockage of blood flow. This observation is a basis for the speculation that changes in Ca^{++} pump system activity is associated with changes in Ca^{++} in white blood cell cytosol; reduced activity of white blood cells as observed during halothane anaesthesia [35-38]. Therefore, white blood cell activity would depend on the level of active Ca^{++} . Dirnagl et al. and Warnke et al. believe that increased white blood cell activity is associated with hypoperfusion [39,40]. Multiple substances are responsible for increased white blood cell activity in the hypoxic tissue. Leukotrienes and peroxides disrupt glycoprotein complex that adheres to the white blood cell surface. Peroxides also change the behaviour of white blood cells in the microcirculation, by stimulating endothelial release of GMP-140

(Granule Membrane Protein 140) (integral membrane glycoprotein) that causes enhanced adhesion of white blood cells to the vessel wall. TNF and interleukin-1 are responsible for production of a factor that causes increased activation of blood platelets along with stimulation of monocytes/macrophages binding to a specific endothelial receptor. All these events result in enhanced white blood cell adhesion to the vascular endothelium, promoting its destruction, microcirculatory, release of toxic agents (oxygen free radicals), proaggregatory factors produced by white blood cells (PAF), resulting in slowing of blood flow in the microcirculation and migration of white blood cells to extravascular tissue [39,41,42].

Blais and Akata believe that in a certain situation, inhaled anaesthetics, including halothane and sevoflurane, may be responsible for vascular spasm and increased permeability through release of oxygen free radicals [2,43,44]. Both anaesthetics could result in such phenomenon through their effect on endothelial guanylate cyclase [45-48]. Akata et al. used 1 MAC for halothane and sevoflurane in their studies but did not find any interference with endothelial guanylate cyclase [43]. Mechanism of this phenomenon is not entirely clear, however vascular contraction in the postcapillary venule seems to be triggered by oxygen free radicals in the presence of inhaled anaesthetics. It is also triggered by xanthine oxidase, flavonoid enzymes and electron transport, resulting in formation of specific, vasoconstrictive prostanoids [5,49-51]. Moreover, protective effects of halogenated anaesthetics were more pronounced in arterial than in venous microcirculation: first, vein walls are almost three-fold less sensitive to inhaled anaesthetics than arterial walls [8], second, protective effect of any anesthetic on metabolic abnormalities in the venous circulation must be less pronounced since hypoxemia is a physiological phenomenon there. Park et al. clearly confirmed that isoflurane had similar mechanism of action to that of sevoflurane and halothane and was responsible for concentration-dependent release of oxygen free radicals [52]. On the contrary, Shayevitz et al. claim that although oxygen free radicals are responsible for injury of the vascular endothelium in the presence of halothane and isoflurane, these anaesthetics are not responsible for the release of reactive oxygen species or their enhanced production, but simply they cause endothelial hyperresponsiveness to these compounds [53-55].

Drinagl et al. studied brain ischemia related

to hypoperfusion and subsequent reperfusion. The authors observed that during hypoperfusion, in particular at its onset, white blood cells form plugs that block the blood flow in microcirculation. However, they were not directly responsible for postischemic hypoperfusion of cerebral cortex, but other, local agent (unidentified by the authors) played dominant role [39]. Structure of brain microcirculation is significantly different from anatomical and physiological characteristics of the peripheral microcirculation [56]. In vitro studies proved that undisturbed white blood cells migrated through an artificial blood brain barrier and did not damage it, while permeability to plasma proteins was unchanged. These studies suggest that this phenomenon may be associated with white blood cell deformation during CNS ischemia that facilitates their migration from the vascular lumen, while in peripheral circulation they migrate outside the vascular lumen only after the endothelium has been damaged [57,58]. On one hand, this phenomenon may be beneficial, since it facilitates reperfusion. However, on the other hand damage processes related to increased white blood cell activity, take place directly in tissues. However, it seems that reduced white blood cell activity in the lumen of the postcapillary venule (white blood cells rolling and adhering to its endothelium) under halothane and sevoflurane anaesthesia favors better capillary perfusion and does not disturb important metabolic processes. If most of toxic mediators related to white blood cell aggregation in the tissues would cause its destruction, these processes must be triggered by intracellular Ca⁺⁺, while halothane and sevoflurane are inhibitors of the latter [38]. These considerations may have practical implications for the selection of anaesthetics for complex and prolonged surgical procedures involving reconstruction of a complex transplants composed of several tissues, including face transplantation.

Conflict of interest

Brak/None

Correspondence address:

✉ Krzysztof Kusza

Chair and Department of Anaesthesiology and Intensive Therapy

Poznan University of Medical Sciences

14, Św. Marii Magdaleny, 61-861 Poznan, Poland

☎ (+48 61) 852 66 87

✉ anestezjologia@ump.edu.pl

References

1. Bagge U, Amudson B, Lauritzen C. White blood cell deformability and plugging of skeletal muscle capillaries in hemorrhagic shock. *Acta Physiol Scand.* 1980;180:159-63.
2. Bloch EH. A quantitative study of the hemodynamics in the living microvascular system. *Am J Anat.* 1962;110:125-53.
3. Branemark PI, Shepro D, Fulton GP. *Rheological Aspects of low flow states. Microcirculation as Related to Shock.* New York: Academic Press; 1968.
4. Chen TC, Skalak R. Stokes flow in a cylindrical tube containing a line of spheroidal particles. *Appl Sci Res.* 1970;22:403-41.
5. Chien S, Sung KP, Schmid-Schonbein GW, Skalak R, Schmalzer EA, Usami S. Rheology of leukocytes. *Ann N Y Acad Sci.* 1989;14:333-47.
6. Dirnagl U, Niwa K, Sixt G, Villringer A. Cortical hypoperfusion after global forebrain ischemia in rats is not caused by microvascular leukocyte plugging. *Stroke.* 1994;25:1028-38.
7. Duncan PG, Cullen BF. Neutrophil chemotaxis and anaesthesia. *Br J Anaesth.* 1977;49:345-9.
8. Nicoll RA, Webb RL. Blood circulation in the subcutaneous tissue of the living bat's wing. *Ann N Y Acad Sci.* 1946;46:697-711
9. Schmid-Schonbein GW, Skalak R, Usami S, Chien S. Cell distribution in capillary networks. *Microvasc Res.* 1980;19:18-44.
10. Nolte D, Bayer M, Lehr HA, Becker M, Krombach F, Kreimeier U, et al. Attenuation of postischemic microvascular disturbances in striated muscle by hyperosmolar saline dextran. *Am J Physiol.* 1992;263:H1411-6.
11. Pardi R, Inverardi L, Bender JR. Regulatory mechanism in leukocyte adhesion: Flexible receptors for sophisticated travelers. *Immunol Today.* 1992;13:224-30.
12. Schmid-Schonbein GW, Zweifach BW, Kovalchek S. The application of stereological principles to morphometry of the microcirculation in different tissues. *Microvasc Res.* 14:303-17.
13. Schmidt-Schonbein GW. Capillary plugging by granulocytes and the no-reflow phenomenon in the microcirculation. *Fed Proc.* 1987;46:2397-401.
14. Siemionow M. *Microcirculatory hemodynamics on muscle flap in vivo.* Thesis for the habilitation. Poznań: AM; 1999. 2,3:42-6.
15. Skalak TC, Schmid-Schonbein GW. The microvasculature in skeletal muscle. IV. A model of the capillary network. *Microvasc Res.* 1986;32:333-47.
16. Fenton BM, Wilson D, Cokelet GR. Analysis of the effects of measured white blood cell entrance times on hemodynamics in computer model of a microvascular bed. *Pflugers Arch.* 1985;403:396-401.
17. Needham D, Hochmuth R. Rapid entry of passive neutrophils into micropipetes. *FASEB J.* 1989;3:693.
18. Amudson B, Jennische E, Haljamae H. Correlative analysis of microcirculatory and cellular metabolic events in skeletal muscle during hemorrhagic shock. *Acta Physiol Scand.* 1980;108:147-58.
19. House SD, Lipowsky H. Leukocyte-endothelium adhesion: Microhemodynamics in mesentery of the cat. *Microvasc Res.* 1987;34:363-79.
20. Mayrovitz HN, Kang SJ, Hescovici B, Sampsell RN. Leukocyte adherence initiation in skeletal muscle capillaries and venules. *Microvasc Res.* 1987;33:22-34.
21. Mayrovitz HN, Rubin R. Leukocyte distribution to arteriolar branches: dependence on blood flow. *Microvasc Res.* 1985;29:282-94.
22. Zweifach BW. (1939) The character and distribution of blood capillaries. *Anat Rec.* 1939;73:475.
23. Zweifach BW, Metz DB. Regional differences in response of terminal vascular bed to vasoactive agents. *Am J Physiol.* 1955;182:55.
24. Gaetgens P, Ley K, Pries AR, Müller R. Mutual interaction between leukocytes and microvascular blood flow. *Prog Appl Microcirc.* 1985;7:15-28.
25. Skalak R. Theoretical modeling of fluid transport through endothelial junctions. *Adv Exp Med Biol.* 1998;242:9-16.
26. Warnke KC, Skalak TC. Leukocyte plugging in vivo in skeletal muscle arteriolar trees. *Am J Physiol.* 1992;262:H1149-55.
27. Morisaki H, Aoyama Y, Shimada M, Ochiai R, Takeda J. Leucocyte distribution during sevoflurane anaesthesia. *Br J Anaesth.* 1998;80:502-3.
28. Warnke KC, Skalak TC. The effects of leukocytes on blood flow in a model skeletal muscle capillary network. *Microvasc Res.* 1990;40:118-36.
29. Gaetgens P. Pathways and interaction of white cells in the microcirculation. *Prog Appl Microcirc.* 1987;12:51-66.
30. Cherry PD, Furchgott RF, Zawadzki JV, Jothianandan D. Role of endothelial cells in relaxation of isolated arteries by bradykinin. *Proc Natl Acad Sci USA* 1982;79:2106-10.
31. Anderson GL, Acland RD, Siemionow M. Vascular isolation of the rat cremaster muscle. *Microvasc Res.* 1988;36:56-63.
32. [195] – odniesienie w wersji oryginalnej; brak pozycji w piśmiennictwie
33. Acland RD, Anderson GL, Siemionow M. Direct in vivo observations of embolic events in microcirculation distal to small-vessel anastomosis. *Plast Reconstr Surg.* 1989;84:280-9.
34. Covino B, Fozzard H, Rehder K, et al. Effects of anesthesia. Cardiovascular effects of general anesthesia. American Physiological Society, Bethesda. 1985;12:149-77.
35. Daniel R, Taylor GI. Distant transfer of an island flap by microvascular anastomoses. *Plas Reconstr Surg.* 1973;52:111-7.
36. Harris PD, Longnecker DE. Significance of precapillary sphincter activity for microcirculatory function. *Microvasc Res.* 1971;3:385-92.
37. Hayes J, Havaleshko D, Plachinta V, Rich GF. Isoflurane pretreatment supports hemodynamics and leukocyte rolling velocities in rat mesentery during lipopolysaccharide-induced inflammation. *Anesth Analg.* 2004;98:999-1006.

38. Den Boer Mo, Van Woerkens LJ, Somers JA, Duncker DJ, Lachmann B, Saxena PR, et al. On the preservation and regulation of vascular tone in arteriovenous anastomoses during anesthesia. *J Appl Physiol.* 1993;76(2):782-9.
39. Folkow B. Role of the nervous system in the control of vascular tone. *Circulation.* 1960;XXI:760-8.
40. Folkow B. Transmural pressure and vascular tone some aspects of an old controversy. *Arch Int Pharmacodyn Ther.* 1962;139:455-69.
41. Yano H, Takaori M. The microcirculation during enflurane and isoflurane anaesthesia in dogs. *Can J Anaesth.* 1994;41:149-55.
42. Yano H, Takaori M. The Microcirculation during enflurane and isoflurane anaesthesia in dogs. *Can J Anaesth* 1994;41:149-55.
43. Akata T, Nakashima M, Kodama K, Boyle WA, Takahashi S. Effects of volatile anesthetics on acetylcholine induced relaxation in the rabbit mesenteric resistance artery. *Anesthesiology.* 1995;82:188-204.
44. Bouskela E, Gramp W. Spontaneous vasomotion in hamster cheek pouch arterioles in varying experimental conditions. *Am J Physiol.* 1992;262:H478-85.
45. Engler RL, Schmid-Schonbein GW, Pavelec RC. Leukocyte capillary plugging in myocardial ischemia and reperfusion in the dog. *Amer J Pathol.* 1983;111:98-111.
46. Hershey SG, Zweifach BW, Rovenstine EA. Effects of depth anesthesia on behaviour of peripheral vascular bed. *Anesthesiology.* 1953;14:245-54.
47. Hsu P, Zuckerman S, Mirro R, Armstead WM, Leffer CW. Effects of ischemia/reperfusion on brain tissue prostanoids and leukotriens in newborn pigs. *Prostaglandins.* 1991;42:557-69.
48. Mellander S. Comparative studies on the adrenergic neuro-hormonal control of resistance and capacitance vessels in the cat. *Acta Physiol Scand.* 1960;50:176-8.
49. Chen G, Yamamoto Y, Miwa K, Suzuki H. Hyperpolarization of arterial smooth muscle induced by endothelial humoral substances. *Am J Physiol.* 1991;260:H1888-92.
50. Erikson E, Myrhage R. Microvascular dimensions and blood flow in skeletal muscle. *Acta Physiol Scand.* 1972;86:211-28.
51. Grant RT. The effects of denervation on skeletal muscle blood vessels (rat cremaster) *J Anat.* 1966;2:305-16.
52. Franks JJ, Horn JL, Janicki PK, Singh G. Halothane, isoflurane, xenon, and nitrous oxide inhibit calcium ATP ase pump activity in rat brain synaptic plasma membranes. *Anesthesiology.* 1995;82:108-17.
53. Franks JJ, Horn JL, Janicki PK, Singh G. Stable inhibition of brain synaptic plasma membrane calcium ATP ase in rats anesthetized with halothane. *Anesthesiology.* 1995;82:118-28.
54. Fujii K, Heistad DD, Faraci FM. Vasomotion of basilar arteries in vivo. *Am J Physiol.* 1990;258(6 Pt 2):H1829-34.
55. Fujihara H, Fukuda S, Fujiwara N, Shimoji K. The effects of halothane on arginine-vasopressin-induced Ca⁺⁺ mobilization from the intracellular space in single cultured smooth muscle cells of rat aorta. *Anest Analg.* 1996;83:584-90
56. Furchgott RF, Zawadzki JV. The obligatory role of endothelial cells in the relaxation of arterial smooth muscle by acetylcholine. *Nature.* 1990;288:373-6.
57. Furchgott RF, Jothianandan D. Endothelium dependent and independent vasodilatation involving cyclic GMP. Relaxation induced by nitric oxide, carbon monoxide and light. *Blood Vessels.* 1991;28:52-61.
58. Furchgott RF, Vanhoutte PM. Endothelium-derived relaxing and contracting factors. *FASEB J.* 1989;3:2007-17.

Regulamin ogłaszania prac w kwartalniku *Anestezjologia i Ratownictwo*

Kwartalnik *Anestezjologia i Ratownictwo* publikuje prace dotyczące zagadnień anestezjologii i ratownictwa, a także szeroko rozumianego pogranicza, związanego również z innymi dyscyplinami naukowymi, w których uwzględniono aspekty diagnostyki, terapii oraz profilaktyki chorób.

Prace należy nadsyłać drogą elektroniczną na adres:

redakcja@akademiamedycyny.pl

Przepisy redakcyjne:

1. Do druku przyjmowane są prace w języku polskim lub angielskim.
2. Prace o charakterze doświadczalnym muszą być w zgodzie z wymogami Konferencji Helsińskiej. Autorzy muszą uzyskać pisemną zgodę osób badanych, po wcześniejszym poinformowaniu ich o przebiegu badań i o ewentualnych szkodliwościach z nich wynikających. Prace, których przedmiotem jest człowiek, mogą być wykonywane i publikowane tylko za zgodą Komisji Bioetycznej i nie mogą ujawniać ich danych osobowych bez załączenia ich pisemnej zgody.
3. Nadesłanie pracy do druku jest jednoznaczne ze stwierdzeniem, że praca nie została zgłoszona do innego czasopisma.
4. Manuskrypty niekompletne lub przygotowane w stylu niezgodnym z zasadami podanymi niżej redakcja odsyła Autorom bez oceny merytorycznej. Artykuły przygotowane zgodnie z wymogami poniżej zostają zarejestrowane i przekazane do oceny niezależnym recenzentom. Akceptacja pracy do druku odbywa się na podstawie pozytywnych opinii recenzentów.
5. Data złożenia pracy w Redakcji, jak również data jej przyjęcia do druku, są umieszczone na początku drukowanej pracy.
6. Prace są recenzowane poufnie i anonimowo (podwójna ślepa próba) przez niezależnych Recenzentów z grona ekspertów w danej dziedzinie.
7. Redakcja zapoznaje Autorów z tekstem recenzji, bez ujawnienia nazwisk recenzentów.
8. Recenzent może uznać pracę za:
 - nadającą się do druku bez dokonania poprawek,
 - nadającą się do druku po dokonaniu poprawek według wskazówek Recenzenta, bez konieczności ponownej recenzji,
 - nadającą się do druku po dokonaniu poprawek według wskazówek Recenzenta i po ponownej recenzji pracy,
 - nienadającą się do druku.
9. Prace wymagające korekty zostaną niezwłocznie przesłane Autorom wraz z uwagami Recenzenta i Redakcji.
10. W przypadku zakwalifikowania pracy do druku Autorzy zostaną o tym fakcie poinformowani e-mailowo lub telefonicznie.
11. Korekty, w formie elektronicznej, należy zwrócić w terminie do 7 od daty wysłania z Redakcji. W wyjątkowych wypadkach Redakcja może przedłużyć termin zwrotu korekty po wcześniejszym uzgodnieniu tego faktu z jej przedstawicielem.
12. Prace niezakwalifikowane do druku zostaną przez Redakcję zniszczone.
13. Redakcja Naukowa zastrzega sobie prawo do dokonywania koniecznych poprawek i skrótów bez porozumienia z Autorami.
14. Prace zgłaszane do druku należy przesłać w formie elektronicznej (e-mail).
15. Objętość prac oryginalnych nie może przekraczać 10 stron, 12 stron dla poglądowych i 8 stron dla prac kazuistycznych. Na jednej stronie nie można umieścić więcej niż 1800 znaków wraz ze spacjami.
16. Wersja elektroniczna pracy powinna być dostarczona e-mailem. Materiał ilustracyjny należy przygotować w formacie TIFF dla materiałów zdjęciowych i skanowanych, a dla grafiki wektorowej w programach Corel Draw do wersji 10 lub Adobe Illustrator do wersji 10, dla wykresów i diagramów MS Excel lub Word.
17. Tekst oraz materiał ilustracyjny powinny być zapisane w oddzielnych plikach np. nazwa-tekst.doc, nazwa-tabela.doc.
18. Obowiązkowy układ pracy:
Strona tytułowa:
 - tytuł artykułu w języku polskim i angielskim,
 - imiona i nazwiska Autorów,

- pełna nazwa instytucji,
- słowa kluczowe w języku polskim/angielskim (do 6 słów),
- pełny adres korespondencyjny jednego z Autorów,
- streszczenie pracy w języku polskim (200-250 słów w przypadku prac oryginalnych oraz 100-150 w przypadku prac poglądowych i opisów przypadków klinicznych; struktura jak w tekście głównym),
- streszczenie pracy w języku angielskim (200-250 słów w przypadku prac oryginalnych oraz 100-150 w przypadku prac poglądowych oraz opisów przypadków klinicznych; struktura jak w tekście głównym),
- należy wyszczególnić wszystkie źródła finansowania wykonanej pracy naukowej.

Układ tekstu głównego (w przyjętym układzie dla prac oryginalnych):

- Wstęp,
 - Cel pracy,
 - Materiał i metody,
 - Wyniki,
 - Omówienie,
 - Wnioski,
 - Podziękowania,
 - Spis piśmiennictwa.
19. Ryciny, tabele, wykresy i fotografie do włączenia w tekst należy nadsyłać oddzielnie, poza tekstem, w którym muszą być zacytowane. Wszystkie powinny być ponumerowane zgodnie z kolejnością występowania w pracy i opisanie w języku polskim i angielskim (tabele - numeracja cyframi rzymskimi; ryciny, wykresy i fotografie – numeracja cyframi arabskimi).
 20. Spis piśmiennictwa powinien ograniczyć się do niezbędnych pozycji cytowanych w pracy, w przypadku prac oryginalnych do 20, a poglądowych do 40 pozycji. Piśmiennictwo należy przytoczyć w kolejności cytowań w tekście. Każdą publikację podaje się w tekście za pomocą cyfry arabskiej w nawiasie kwadratowym. Cytowanie w spisie piśmiennictwa powinno mieć następujący układ:
 - dla czasopisma: nazwiska autorów z inicjałami imion, oddzielone przecinkami, zakończone kropką (jeśli liczba autorów cytowanej pracy przekracza sześć osób, to pozostałych należy zaznaczyć skrótem: „i wsp.”), tytuł artykułu, kropka, nazwa czasopisma przytoczona w skrócie wg Index Medicum, kropka, spacja, rok, średnik, tom, dwukropek, pierwsza i ostatnia strona – (np.: 1. Jakobsson U, Hallberg IR. Loneliness, fear and quality of life among elderly in Sweden: a tender perspective. *Aging Clin Exp Res.* 2005;17(6):494-501.
 - dla tytułu rozdziału z książki: nazwiska autorów z inicjałami imion, oddzielone przecinkami, zakończone kropką, tytuł rozdziału, kropka, W: nazwiska redaktorów (red.), kropka, tytuł książki, kropka, miejscowość, dwukropek, spacja, wydawca, średnik, spacja, rok wydania, kropka lub – w przypadku wybranych stron - dwukropek, spacja, pierwsza i ostatnia strona – np.: Wojszel ZB, Bień B. Wielkie problemy geriatryczne – rola zespołu terapeutycznego w opiece nad pacjentem. W: Kędziora-Kornatowska K, Muszałik M (red.). *Kompendium pielęgnowania pacjentów w starszym wieku. Podręcznik dla studentów i absolwentów kierunku pielęgniarstwo.* Lublin: Wydawnictwo Czelej; 2007. str. 97-114.
- Dopuszcza się również cytowania ze źródeł elektronicznych.
21. Praca powinna być zredagowana możliwie krótko, bez zarzutu pod względem stylistycznym, zgodnie z obowiązującą pisownią.
 22. Należy używać międzynarodowych (zgodnie z zasadami polszczyzny) nazw leków. Dopuszcza się podawanie nazw handlowych w nawiasach.
 23. Skróty powinny być wyjaśnione w tekście w miejscu, w którym się pojawiają po raz pierwszy.
 24. Wraz z pracą należy złożyć Deklarację Konflikty Interesów oraz List Przewodni z oświadczeniem Autorów, że praca nie została i nie zostanie złożona do druku w innym czasopiśmie oraz że nie zachodzą zjawiska: „guest authorship” i „ghostwriting”.
 25. Pierwszy autor/autor do korespondencji otrzymują bezpłatnie 1 egzemplarz czasopisma z wydrukowanym artykułem.
 26. Nie przewiduje się honorariów autorskich
 27. Prawa autorskie: Maszynopis zakwalifikowany do druku w kwartalniku staje się własnością Wydawnictwa Akademia Medycyny. Wydawca nabywa na zasadzie wyłączności ogół praw autorskich do wydrukowanych prac (w tym prawo do wydawania drukiem, na nośnikach elektronicznych i innych oraz w Internecie). Bez zgody wydawcy dopuszcza się jedynie drukowanie streszczeń.
 28. Redakcja *Anestezjologii i Ratownictwa* posiada własną stronę internetową, na której zamieszczane są streszczenia drukowanych prac, jak również istotne wiadomości.

Submission Manuscript Guidelines:

The quarterly journal *Anestezjologia i Ratownictwo / Anaesthesiology and Rescue Medicine* publishes articles which cover key issues and current trends in geriatric medicine, as well as it also presents the broadest disciplines that focus on any aspect of the diagnosis, therapy and the prevention of the ageing related diseases.

Please submit your papers electronically at:

redakcja@akademiamedycyny.pl

Instructions for authors

1. Only papers written in Polish or English are accepted.
2. In scientific investigations involving human subjects, experiments should be performed in accordance with the ethical standards formulated in the Helsinki Declaration. Informed consent for the research must be obtained from all participants and all clinical investigations. For papers involving human subjects, adequate documentation should be provided to certify that appropriate ethical safeguards and protocols have been followed according to the responsible Bioethical Committee on human experimentation (institutional or regional). Names should not be published in written descriptions, photographs, sonograms, CT scans, etc., nor should pedigrees, unless information is essential for scientific purposes and a patient (or a parent, or a guardian) has given their written informed consent for publication.
3. Manuscripts are received with the explicit understanding that they are not under simultaneous consideration by any other publication. Submission of an article implies that the work described has not been published previously.
4. Manuscripts that are incomplete or whose style does not follow the below guidelines shall be returned to the Authors without being evaluated. Articles following the below guidelines shall be registered and sent to independent reviewers to be evaluated. A paper shall be accepted for publication based on reviewers' positive opinions.
5. The dates of submitting and acceptance for publication are labeled at the end of each manuscript.
6. Submitted manuscripts are anonymously reviewed by two impartial experts to determine their originality ("double-blind review"), scientific merit, and significance to the field.
7. Reviewers shall remain anonymous, but their comments will be available to authors.
8. There are several types of decision possible:
 - accept the manuscript as submitted,
 - accept it with revision,
 - accept it and invite the authors to revise the manuscript before a final decision is reached,
 - accept it with encouragement to resubmit it after extensive revision,
 - outright rejection.
9. Page proofs with reviewer's remarks will be sent to corresponding author for examination and corrections.
10. Information about accepting the manuscript for publication will be sent to the corresponding author.
11. Corrected proofs should be returned to the Editor within seven days of posting by the Editor. Authors are responsible for obtaining the Editor's permission for any changes in the time for returning proofs.
12. When submitted manuscripts are not accepted for publication, they will be destroyed according to the Editorial office's schedule.
13. The Editors reserve the right to make corrections in style and nomenclature without Authors' permission.
14. Authors should return the final, revised manuscript by e-mail: redakcja@akademiamedycyny.pl
15. Manuscripts of original papers should not exceed 10 pages, review articles – 12 pages, case reports – 8 pages. One page is generally limited to 1800 characters including spacing.
16. The electronic version of the text should be submitted as MS Word 98 or above. All illustration and scan files should be in the TIFF format. For vector graphics, the digital formats of Adobe Illustrator for version 10 and Corel Draw for version 10 are accepted; for graphs and diagrams - MS Excel or MS Word.
17. The text and figures must be uploaded as separate files. Files should be named with the corresponding Author's surname and "text.doc", "fig. 1.doc", "fig. 2.doc" name and extension formats, etc.
18. The paper should be laid out as follows:
Provide the following data on the title page (in the order given):
 - the article's title,
 - Authors' names,

- institutional affiliations, the name of department(s) and institution(s) to which the work should be attributed,
- keywords (maximum of 6 keywords),
- full postal address of the corresponding Author,
- an abstract in English (maximum length of 200-250 words in case of original works and 100-150 in case of review articles or clinical cases, an abstract should state briefly the purpose of the research, principal results and major conclusions.

Arrangement of the article (for original papers):

- Introduction,
 - Purpose of the work,
 - Material and methods,
 - Results,
 - Discussion,
 - Conclusions,
 - Acknowledgements,
 - References,
 - Source(s) of support in the form of grants, equipment, drugs, or all of these.
19. Tables, illustrations, vector graphics and photographs should be prepared and submitted on separate pages. All figures should be numbered in the order of their citation in the text and legends should be in Polish and English (tables - Roman numerals; illustrations, vector graphics and photographs - Arabic numerals).
 20. References should exceed in number, and should in general be limited according the paper type: for original papers – up to 20, for review articles – up to 40 items. They must be numbered in their order of appearance in the text. References should be identified in the text, tables, and legends by Arabic numerals in square brackets. It is allowed to use the following style for the references list: surname and initials of all authors separated by a comma, followed by a full stop, then the article title (or the book title), a full stop, the name of the journal should be abbreviated according to the style used in the Index Medicus, the year, a semicolon, the volume number, a colon, the first and the last page (for books: the city, a colon, the publisher, a semicolon, the year, a colon, pages). When there are more than six authors, only the first six authors are listed, followed by “et al.”, i.e.: 1. Jakobsson U, Hallberg IR. Loneliness, Fear and quality of life among the elderly in Sweden: a tender perspective. *Aging Clin Exp Res.* 2005;17(6):494-501. 2. Wojszel ZB, Bień B. Wielkie problemy geriatryczne – rola zespołu terapeutycznego w opiece nad pacjentem. In: Kędziora-Kornatowska K, Muszalik M (ed.). *Kompendium pielęgnowania pacjentów w starszym wieku. Podręcznik dla studentów i absolwentów kierunku pielęgniarstwo.* Lublin: Wydawnictwo Czelej; 2007. p.: 97-114.
- Quoting from electronic sources is accepted.
21. A paper ought to be brief and observe general style and spelling rules.
 22. International generic rather than trade names of drugs should be used. Trade or manufacturers’ names should only be used in brackets.
 23. All abbreviations should be spelt out in full the first time they are used.
 24. A paper submitted for publication should be accompanied by a Declaration of ‘Conflict of Interest’ and a ‘Cover Letter’ with a statement by the Author(s) confirming that the paper has not been and will not be published elsewhere and that there is no instance of misconduct (“ghostwriting” and “guest authorship”).
 25. First Author/corresponding Author shall receive 1 copy of the issue in which the article shall be published. They shall be sent to the author(s) free of charge.
 26. No remuneration shall be paid for publication.
 27. Copyrights. Submission of an article for publication implies a transfer of the copyright from the Author to the publisher upon acceptance. Accepted papers shall become permanent property of the Wydawnictwo Akademia Medycyny and may not be reproduced without the written consent from the publisher. The publisher reserves the copyright (including printing, electronic version such as CDs, the Internet and others). Only abstracts can be published elsewhere without written permission from the publisher.
 28. A website shall be launched by the *Anaesthesiology and Rescue Medicine*, which shall feature abstracts of printed papers as well as other relevant information.